JOURNAL OF GEOSPACE SCIENCE

Vol. 6 No. 2 (2026)

E-ISSN : 2413-6093
https://jogss.org

Analysis of Ionospheric Total Electron Content Variations During Solar and Geomagnetic Disturbances

Dr. Elena Maren Solvik
Department of Space Physics, North Atlantic Institute for Geospace Studies, Bergen, Norway

ABSTRACT

Ionospheric total electron content provides an integrated measure of the plasma through which satellite radio signals propagate. Its variability reflects solar radiation, magnetospheric energy input, neutral atmospheric dynamics, and redistribution of charged particles. This review examines how solar and geomagnetic disturbances modify total electron content and how those modifications should be measured and interpreted. The discussion connects dual frequency satellite observations with ionospheric production, recombination, electric fields, thermospheric circulation, and regional differences in storm response. Particular attention is given to the separation of rapid solar flare signatures from delayed geomagnetic effects, the selection of quiet reference conditions, and uncertainty introduced by instrumental biases and mapping assumptions. A conceptual graph illustrates positive and negative departures from a quiet baseline using explicitly synthetic values; no new observational dataset or event statistics are reported. The synthesis indicates that total electron content alone cannot uniquely identify a disturbance mechanism or quantify scintillation risk. Reliable attribution requires complementary measurements, consistent temporal alignment, and assessment of both spatial gradients and background variability. The article proposes a reproducible analytical framework and explains its relevance to satellite navigation, communication reliability, and operational space weather monitoring.

KEYWORDS

ionospheric total electron content, geomagnetic storms, solar flares, GNSS observations, space weather

1. Introduction

Total electron content, commonly abbreviated TEC, is the number of free electrons integrated along a specified propagation path through near Earth plasma. It connects ionospheric physics with radio engineering because dispersive propagation changes the apparent distance measured by a satellite receiver. Variations that appear modest in an atmospheric context may matter to precise navigation, timing, surveying, and communication systems. Understanding those variations therefore requires both a physical explanation of the plasma and an explicit account of how the measurement is obtained.Solar and geomagnetic disturbances influence TEC through different pathways. Increased solar ionizing radiation can modify electron production on the illuminated hemisphere within a short interval. Solar

wind disturbances affect the magnetosphere and subsequently alter electric fields, particle precipitation, heating, neutral winds, and chemical loss. These processes have different delays and spatial footprints. A TEC enhancement observed during a geomagnetic storm may consequently arise from redistribution, additional production, reduced loss, or several mechanisms acting together. A decrease may likewise reflect chemical depletion, transport, or changing contributions from the topside ionosphere.The distinction matters because TEC is an integral rather than an altitude profile. Two columns can contain similar electron numbers while differing substantially in peak density, layer height, thickness, and irregularity structure. Ground based satellite measurements also sample changing ray paths. Apparent temporal changes can partly arise when a moving satellite observes different regions. The physical and observational meanings of a variation must therefore be evaluated together before a causal interpretation is assigned.This article develops a narrative review and an analytical framework rather than presenting original experimental results. It draws on established ionospheric physics, published storm studies, satellite measurement methods, and propagation research. The twelve references were selected for their relevance to these themes; the selection is not a systematic literature search or a claim of exhaustive coverage. No receiver archive, event catalogue, or newly measured solar wind series was processed. The numerical example is synthetic and serves only to explain analysis choices.The central argument is that robust disturbance analysis depends on comparing calibrated observations with an appropriate baseline while retaining information about location, local time, uncertainty, and scale. The following nine sections examine physical foundations, external drivers, observations, processing, regional responses, disturbance metrics, illustrative interpretation, technological consequences, and future research. Together they explain what TEC measurements can establish and where additional evidence is necessary.

Physical Foundations of Total Electron Content:

TEC is expressed as the path integral $TEC = \int n_e \, ds$, where n_e denotes electron number density and ds is a distance element along the ray. One TEC unit, or TECU, equals 10^{16} electrons per square metre. Slant TEC describes the actual satellite to receiver line of sight, whereas vertical TEC represents an equivalent overhead column. These quantities should never be compared without acknowledging their different geometry. The distinction becomes especially consequential at low satellite elevations, where the ray crosses a longer and more horizontally extensive plasma region.Electron density evolves through production, loss, and transport. Solar extreme ultraviolet radiation produces much of the dayside ionization, while chemical reactions remove electrons at rates that depend on composition and temperature. Electric fields, pressure gradients, gravity, and collisions influence motion. A continuity statement can be written as $\partial n_e/\partial t = P - L - \nabla\cdot(n_e v)$, with P representing production, L loss, and v plasma velocity. Integrating this balance helps explain why a change in TEC cannot automatically be interpreted as a proportional change in solar radiation.An uplifted plasma layer can experience slower chemical loss because molecular neutral densities generally decline with altitude. TEC may then increase even without a comparable increase in ionizing flux. Conversely, transport into a region with more efficient loss can reduce the integrated column. The relative importance of these effects changes with latitude, season, local time, and storm phase. The plasma and electrodynamic framework described by Kelley (2009) provides the basis for interpreting such competing influences.Ground to navigation satellite TEC includes electrons above the ionospheric density peak and can contain a plasmaspheric contribution. Thus, a persistent change may involve storage and exchange beyond the altitude range most directly sampled by an ionosonde. Schunk and Nagy (2009) describe the interacting chemical and plasma processes needed to interpret this vertical structure. An observational study should state the propagation endpoints and avoid treating every reported TEC product as an identical atmospheric column.The normal daily cycle provides essential context. TEC commonly grows after sunrise, reaches an afternoon maximum, and decreases overnight, although regional electrodynamics complicate this pattern. Seasonal and solar cycle changes alter both the baseline and the available plasma reservoir. A disturbed afternoon value can exceed a quiet night value while still representing a negative anomaly

relative to the appropriate afternoon reference. The sign of a storm response therefore belongs to a defined comparison, not to the absolute electron content alone.

Solar Radiation and Geomagnetic Drivers:

Solar flares and geomagnetic storms should be separated in both the event description and the analysis. Flare radiation reaches the sunlit atmosphere directly and can generate a rapid increase in ionization. The timing and amplitude of the TEC response depend on spectral irradiance, solar zenith angle, atmospheric state, and the distribution of electrons with height. A flare classification based on one radiation band does not by itself specify the response of the integrated column or every radio propagation effect.Afraimovich (2000) introduced coherent use of a global GPS network to detect ionospheric flare responses. The important analytical principle is that a physically coordinated dayside response can become clearer when many independent ray paths are examined together. For a new study, coherence should be tested against instrument artefacts, satellite geometry, and the expected illumination pattern. A sudden change on one link is insufficient evidence of a global solar signature.Geomagnetic disturbances develop when solar wind energy and momentum are transferred into the magnetosphere and upper atmosphere. The interplanetary magnetic field orientation, solar wind speed, density, and dynamic pressure help characterize this forcing. Southward magnetic field intervals can favor magnetic reconnection, although no single upstream variable fully describes the ionospheric response. Propagation delays from an upstream monitor must be handled explicitly before its time series is compared with ground observations.Fast changes in magnetospheric forcing can modify ionospheric electric fields and redistribute plasma. Slower heating and circulation changes can alter composition and sustain effects beyond the strongest magnetic disturbance. Fuller Rowell et al. (1994) examined the coupled thermosphere and ionosphere response, showing why neutral atmospheric dynamics belong in a storm interpretation. An early enhancement and a later depletion can therefore reflect different stages of the same coupled disturbance rather than inconsistent observations.Geomagnetic indices provide complementary context. Dst and SYM H describe broad low latitude magnetic perturbations associated with storm evolution, while Kp summarizes planetary magnetic activity and AE characterizes auroral electrojet activity. Their sampling and spatial meanings differ. Aligning TEC exclusively to the minimum of one index can obscure an earlier local response. Event analysis should retain driver chronology and distinguish shock arrival, changing electric forcing, main phase development, and recovery without assuming that all regions follow the same sequence.

Observation Systems and Complementary Measurements:

Dual frequency Global Navigation Satellite System observations are valuable because the leading ionospheric propagation term depends on inverse frequency squared. Combining measurements at two frequencies separates much of this dispersive contribution from nondispersive effects. Code observations provide an absolute but relatively noisy estimate, while carrier phase observations resolve changes precisely but include ambiguity offsets. A reliable TEC series consequently requires more processing than simply subtracting two recorded satellite ranges.Satellite and receiver differential code biases must be estimated or obtained from suitable calibration products. Their units, sign conventions, reference observables, and time validity need to match the processing software. Mixing incompatible bias conventions can create systematic offsets that resemble geographic differences. Rideout and Coster (2006) describe automated processing for global TEC products, demonstrating the importance of consistent calibration when observations from many receivers are combined.Receiver networks sample the ionosphere unevenly. Dense land networks resolve regional structure more effectively than sparsely observed oceans and polar areas. A smooth global map can therefore contain both strongly constrained regions and locations where interpolation dominates. The availability of several constellations improves sampling but also introduces intersystem and receiver related differences. A map should be assessed together with coverage information rather than assumed to have uniform reliability across its entire domain.Complementary instruments help address the vertical ambiguity of TEC. Ionosondes estimate F region peak characteristics; incoherent scatter radars constrain density, temperature, and velocity

profiles; satellite radio occultation provides altitude dependent information under retrieval assumptions. Airglow imaging and in situ measurements can reveal structures that an integrated map cannot separate. Agreement across instruments is strongest when their observation volumes, times, and sensitivity ranges are explicitly matched.Useful comparisons also include thermospheric composition observations, neutral wind measurements, solar irradiance, and magnetic field records. Each constrains a different link in the causal chain. A TEC increase accompanied by layer uplift supports a transport related interpretation more strongly than TEC alone, but still requires attention to chemical changes and horizontal advection. The purpose of combining instruments is to test competing explanations, not merely to collect several plots that change during the same storm.

Data Processing and Quiet Reference Construction:

A reproducible analysis begins with a clear definition of the event interval, geographical domain, receiver selection, observables, and time standard. Processing should retain raw data provenance, receiver metadata, orbit products, calibration files, and software versions. Times should be expressed consistently in universal time, with local solar time or magnetic local time added where appropriate. These choices allow another investigator to reconstruct which plasma regions were sampled and how the reported anomalies were calculated.Quality control should identify cycle slips, phase discontinuities, multipath, low elevation observations, data gaps, and receiver changes. Carrier phase arcs need separate treatment when ambiguities reset. Interpolation across a gap can smooth a real disturbance or create an apparent transition that was never observed. Removing questionable data is reasonable, but rejection rules should be stated before inspecting the expected storm response and the resulting loss of coverage should be reported.Mapping slant measurements into vertical TEC often uses a thin shell approximation. For elevation E, Earth radius R, and shell height h, a common mapping factor is $M = [1 - (R \cos E/(R + h))^2]^{-1/2}$, with VTEC = STEC/M. This representation is convenient but assumes a simplified vertical distribution. During strong uplift, broad layer changes, or sharp horizontal gradients, its error can vary with geometry. Testing plausible shell heights helps determine whether an inferred anomaly depends on the approximation.Quiet references can be based on selected magnetically quiet days, a rolling median, or a climatological model. Each answers a different question. Selected days may have different solar illumination or radiation conditions; rolling statistics may absorb prolonged disturbances; climatologies may miss ordinary weather variability. The baseline should be constructed for comparable local times and locations, and its spread should be retained rather than reduced to a single smooth curve.Sensitivity analysis is part of the scientific result. An anomaly that persists under several reasonable baselines is more defensible than one produced by a single convenient reference. A useful report gives the event TEC, reference TEC, absolute difference, percentage difference, and uncertainty together. It also states how candidate quiet days were screened. The objective is to show that a disturbance exceeds plausible background and processing variability, rather than to maximize the apparent storm amplitude.

Latitude Local Time and Seasonal Response Patterns:

Low latitude TEC responds strongly to electric field driven vertical motion and the redistribution associated with the equatorial ionization anomaly. Changes in the fountain process can shift anomaly crests as well as alter their amplitude. A fixed receiver near a crest can observe an increase simply because the structure moves toward its sampling region. Measurements spanning several magnetic latitudes are therefore needed to distinguish a migrating feature from a uniform regional increase.Middle latitudes can exhibit storm enhanced density, extended depletions, and traveling disturbances. Their relative contributions depend on background plasma, electric fields, winds, and composition. Buonsanto (1999) reviews the interacting processes behind F region storm behavior. A regional study should examine whether neighboring stations share an anomaly, whether its boundary moves, and whether independent observations support a change in layer height or chemical loss.At high latitudes, precipitation and strong convection produce complicated spatial structure. Polar cap patches, auroral boundaries, and depleted regions can create steep gradients over short distances. A fixed receiver samples different parts of these features as satellites move. Interpreting rapid TEC variation therefore

requires tracking ionospheric pierce points and separating changes caused by plasma evolution from those caused by movement through an existing structure.Local time determines illumination, background density, conductivity, and the direction or effectiveness of several transport processes. Hemispheric and seasonal differences add further complexity. A summer depletion and winter enhancement need not be contradictory because the neutral atmosphere and magnetic geometry differ. Mendillo (2006) places the diversity of TEC storm patterns in historical context and cautions against treating a single regional response as a universal storm template.Published extreme events demonstrate that large positive responses are physically possible without defining the behavior of every disturbance. Mannucci et al. (2005), for example, documented pronounced dayside enhancements during the October 2003 storms. Such observations motivate analysis of electric field driven redistribution, but their amplitudes should not be transferred to an unrelated event. Comparative studies gain explanatory value when they match season, magnetic latitude, local time, solar activity, and preconditioning as carefully as possible.

Disturbance Metrics and Statistical Interpretation:

The absolute anomaly is $\Delta TEC = TEC_{event} - TEC_{reference}$. Its percentage form is $100\Delta TEC/TEC_{reference}$. Both should be reported because a large percentage change at night can correspond to a small absolute column change. Relative metrics become unstable when the reference approaches zero. In that situation, reporting an absolute anomaly with the reference uncertainty is generally more informative than presenting an exaggerated percentage or silently excluding inconvenient observations.Rate of TEC and the rate of TEC index, ROTI, characterize short interval variability along an observation path. ROTI is commonly calculated as the standard deviation of TEC change rates within a specified window. Pi et al. (1997) established the value of network observations for monitoring irregularity related TEC variability. Window length, sampling interval, elevation mask, arc treatment, and units must accompany a reported ROTI value because changes in these settings alter its interpretation.Temporal variability is not identical to spatial structure. A moving pierce point can cross a gradient even when the plasma field evolves slowly. Conversely, a fixed region can change rapidly with little spatial contrast. Detrending choices determine which scales remain in the residual. Filters should therefore be selected according to the physical question and tested on representative signals. A visually prominent oscillation can otherwise be an artefact of a filter edge, interpolation, or abrupt arc termination.Storm comparisons also require statistical independence. Thousands of high cadence samples from a few stations do not provide thousands of independent realizations of a storm mechanism. Serial dependence, shared satellite paths, and regional coherence reduce the effective information content. Event level resampling, station group comparisons, and uncertainty intervals can provide more realistic inference than treating every epoch as an unrelated observation. Sample counts and missing data should remain visible in the reported analysis.Correlation between TEC and a geomagnetic index is useful descriptive evidence but cannot alone establish causation. Both may vary with another driver, and their response delays may change across an event. Lag searches increase the chance of finding a large correlation by chance if many offsets are tested. A defensible analysis specifies hypotheses, evaluates physically plausible delays, compares alternative drivers, and distinguishes predictive association from a demonstrated mechanism. Negative findings should be retained when they constrain an explanation.

Illustrative TEC Variation and Physical Interpretation:

The graph accompanying this section is an original conceptual illustration constructed from synthetic values. It covers an arbitrary interval from twelve hours before a nominal disturbance to forty eight hours afterward. The quiet reference represents a smooth daily cycle. The disturbed curve adds a broad enhancement followed by a depletion. These shapes were selected to explain anomaly analysis; they do not represent a measured storm, a fitted physical simulation, or a statistical average of published events.

For reproducibility, let t be hours relative to the nominal onset. The quiet curve is $Q(t) = 25 + 10\cos[2\pi(t - 2)/24]$ TECU. The illustrated disturbance is $D(t) = Q(t) + 22\exp[-((t - 8)/5)^2] - 12\exp[-((t -$

30)/9)²] TECU. The baseline is always positive, and the disturbance terms create separate enhancement and depletion intervals. The equation contains no estimated coefficients, observational uncertainty, or event specific solar forcing.At t = 8 hours, the quiet value is 25 TECU and the disturbed value is approximately 46.97 TECU, giving an illustrative anomaly of about 21.97 TECU or 87.9%. At t = 30 hours, the quiet value is 30 TECU and the disturbed value is approximately 18 TECU, corresponding to a depletion of 12 TECU or 40%. These calculations demonstrate how identical plotting conventions support both absolute and relative interpretation without implying that these amplitudes are typical.An observed curve with this appearance would invite several hypotheses. The enhancement could reflect uplift, horizontal transport, altered production, or reduced loss. The depletion could reflect composition changes, export of plasma, or reduced replenishment. Testing those possibilities would require measurements absent from this illustration. The nominal onset also does not establish an observational lag: a real analysis would define onset from independent upstream or geomagnetic evidence and account for timing uncertainty.The graph additionally shows why recovery should be evaluated relative to the evolving quiet baseline. A decreasing disturbed curve can approach normal conditions, cross below them, or remain enhanced while the daily background falls. Visual comparison with a horizontal threshold would confuse these outcomes. For measured data, confidence bands, coverage indicators, and separate uncertainty estimates would be needed before the same figure could support event classification or operational decisions.




Figure 1 Synthetic vertical TEC response relative to a daily quiet reference. Values are generated from the equations in the text and are not observations. Shading marks enhancement and depletion relative to the reference.

Consequences for Satellite Navigation and Communications:

The first order ionospheric group delay in metres is approximately I = 40.3 TEC/f² when TEC is in electrons per square metre and frequency f is in hertz. At the GPS L1 frequency of 1.57542 GHz, one TECU corresponds to approximately 0.162 metres of slant range delay. An additional 20 TECU along that ray would therefore add about 3.25 metres before correction. A vertical anomaly must first be related to the actual slant path before this conversion is applied.This calculation represents a propagation contribution, not a direct position error. Receiver location estimates combine many observations whose errors interact with satellite geometry, weighting, correction methods, and estimation algorithms. Klobuchar (1987) developed a compact broadcast correction approach for single frequency users. Its role illustrates why a background model can reduce an expected delay while still leaving event dependent

residuals that require observations or more detailed corrections to characterize.Dual frequency combinations remove most first order ionospheric delay, but they do not guarantee uninterrupted performance. Rapid phase variation, signal fading, cycle slips, tracking failure, and unresolved gradients can still degrade positioning. Coster and Komjathy (2008) describe the connection between space weather and GPS capabilities. The relevant performance assessment should therefore examine ambiguity recovery, continuity, accuracy, and integrity rather than infer service quality solely from a regional TEC maximum.Bulk TEC and scintillation are related through plasma structure but are different observables. Scintillation depends on irregularity scales, propagation geometry, radio frequency, and receiver response. A large smooth TEC increase may generate substantial delay with little rapid fading, whereas structured plasma at moderate background TEC can challenge tracking. Jakowski et al. (2012) discuss monitoring and forecasting ionospheric perturbations for GNSS applications. Their operational perspective supports reporting gradients and variability alongside integrated content.Communication effects also depend on the system. High frequency links respond to layer structure and absorption, while satellite links encounter frequency dependent delay and irregularity related fluctuations. TEC does not replace direct measurements of absorption or channel performance. A practical assessment combines ionospheric products with link frequency, elevation, receiver behavior, and observed service degradation. This permits an evidence based connection between a plasma disturbance and an actual communication consequence.

Forecasting Limitations and Research Priorities:

Forecasting TEC during disturbed conditions remains difficult because external forcing interacts with an atmosphere that already has its own variability. A storm of similar magnetic intensity can produce different regional effects when its onset occurs at another local time or under different background conditions. Useful forecasts should therefore specify their location, horizon, temporal resolution, expected uncertainty, and intended application rather than promise one global disturbance magnitude.Physics based models represent production, chemistry, transport, and coupling, but depend on uncertain drivers and boundary conditions. Empirical models summarize recurring behavior but may smooth unusual events. Data driven systems can capture patterns in large archives, yet may fail when training coverage is unrepresentative. Comparing these approaches requires the same observation definitions and independent evaluation periods. A sophisticated model should be tested against simple persistence and climatology before additional complexity is credited with practical value.Data assimilation can combine models and observations, provided their respective errors are represented realistically. Assimilating biased TEC can improve agreement with one network while degrading the underlying state estimate. Conversely, sparse observations can leave large unobserved regions weakly constrained. Validation should include withheld stations and, where possible, complementary profile observations. This helps distinguish better reproduction of an integral from better representation of the physical ionosphere.Evaluation should emphasize disturbed intervals, regional extremes, gradients, and operational thresholds as well as average error. A system that performs well during numerous quiet hours may still miss the short intervals that matter most to users. Reliability diagrams, event detection statistics, uncertainty coverage, and separate regional scores can reveal those weaknesses. Training and testing by independent events reduces leakage caused by adjacent samples from the same disturbance appearing in both datasets.Further progress depends on consistent metadata, transparent processing, broader receiver coverage, and coordinated observations of plasma and neutral atmosphere. Reusable event analyses should publish code, calibration choices, baseline construction, uncertainty assumptions, and exclusion rules when data permissions allow. The priority is to turn an observed anomaly into a testable physical explanation and an application relevant estimate. That goal is best served by preserving uncertainty rather than assigning a mechanism that the available measurements cannot resolve.

Summary:

Ionospheric TEC variations during solar and geomagnetic disturbances result from interacting changes in production, chemical loss, and plasma transport. Solar flare radiation and magnetospheric forcing

have distinct pathways, while neutral atmospheric circulation and composition can sustain effects through recovery. Latitude, local time, season, and preconditioning determine which influences dominate. Positive and negative anomalies consequently need to be defined relative to a suitable quiet reference rather than inferred from absolute electron content.A reliable analysis requires calibrated dual frequency measurements, explicit geometry, careful quality control, and a baseline with documented variability. Absolute anomalies, percentages, temporal variability, and spatial gradients answer complementary questions. None uniquely identifies the responsible physical mechanism. Independent observations of layer height, density profiles, winds, composition, or electrodynamics strengthen attribution and prevent integrated measurements from being overinterpreted.The synthetic graph demonstrates an enhancement followed by depletion and explains the arithmetic of anomaly estimation. It supplies no evidence about the timing or magnitude of an actual event. For technological applications, TEC is a useful measure of dispersive delay, while scintillation and receiver performance require additional observations. Future studies should combine transparent processing, event independent validation, and uncertainty aware forecasting so that physical interpretation and operational guidance remain proportionate to the evidence.

References:

Afraimovich, E. L. (2000). The GPS global detection of the ionospheric response to solar flares. Radio Science, 35(6), 1417–1424. https://doi.org/10.1029/2000RS002340

Buonsanto, M. J. (1999). Ionospheric storms—A review. Space Science Reviews, 88, 563–601. https://doi.org/10.1023/A:1005107532631

Coster, A., & Komjathy, A. (2008). Space weather and the Global Positioning System. Space Weather, 6, S06D04. https://doi.org/10.1029/2008SW000400

Fuller-Rowell, T. J., Codrescu, M. V., Moffett, R. J., & Quegan, S. (1994). Response of the thermosphere and ionosphere to geomagnetic storms. Journal of Geophysical Research: Space Physics, 99(A3), 3893–3914. https://doi.org/10.1029/93JA02015

Jakowski, N., Béniguel, Y., De Franceschi, G., Hernández-Pajares, M., Jacobsen, K. S., Stanislawska, I., Tomasik, L., Warnant, R., & Wautelet, G. (2012). Monitoring, tracking and forecasting ionospheric perturbations using GNSS techniques. Journal of Space Weather and Space Climate, 2, A22. https://doi.org/10.1051/swsc/2012022

Kelley, M. C. (2009). The Earth's ionosphere: Plasma physics and electrodynamics (2nd ed.). Academic Press.

Klobuchar, J. A. (1987). Ionospheric time-delay algorithm for single-frequency GPS users. IEEE Transactions on Aerospace and Electronic Systems, AES-23(3), 325–331. https://doi.org/10.1109/TAES.1987.310829

Mannucci, A. J., Tsurutani, B. T., Iijima, B. A., Komjathy, A., Saito, A., Gonzalez, W. D., Guarnieri, F. L., Kozyra, J. U., & Skoug, R. (2005). Dayside global ionospheric response to the major interplanetary events of October 29–30, 2003 "Halloween Storms". Geophysical Research Letters, 32, L12S02. https://doi.org/10.1029/2004GL021467

Mendillo, M. (2006). Storms in the ionosphere: Patterns and processes for total electron content. Reviews of Geophysics, 44, RG4001. https://doi.org/10.1029/2005RG000193

Pi, X., Mannucci, A. J., Lindqwister, U. J., & Ho, C. M. (1997). Monitoring of global ionospheric irregularities using the worldwide GPS network. Geophysical Research Letters, 24(18), 2283–2286. https://doi.org/10.1029/97GL02273

Rideout, W., & Coster, A. (2006). Automated GPS processing for global total electron content data. GPS Solutions, 10, 219–228. https://doi.org/10.1007/s10291-006-0029-5

Schunk, R. W., & Nagy, A. F. (2009). Ionospheres: Physics, plasma physics, and chemistry (2nd ed.). Cambridge University Press. https://doi.org/10.1017/CBO9780511635342